

II.H.4 Maximizing Light Utilization Efficiency and Hydrogen Production in Microalgal Cultures

Tasios Melis

University of California, Berkeley
 Dept. of Plant & Microbial Biology
 111 Koshland Hall
 Berkeley, CA 94720-3102
 Phone: (510) 642-8166
 E-mail: melis@berkeley.edu

DOE Managers

HQ: Roxanne Garland
 Phone: (202) 586-7260
 E-mail: Roxanne.Garland@ee.doe.gov
 GO: Katie Randolph
 Phone: (720) 356-1759
 E-mail: Katie.Randolph@go.doe.gov

Contract Number: DE-FG36-05GO15041

Start Date: December 1, 2004
 End Date: November 30, 2011

Approach

- Employ deoxyribonucleic acid (DNA) insertional mutagenesis, screening, biochemical and molecular genetic analyses for the isolation of “truncated Chl antenna size” mutants in the green alga *Chlamydomonas reinhardtii*.
- Clone and characterize the gene(s) that affect the “Chl antenna size” property in *Chlamydomonas reinhardtii*.
- Apply such genes to generate a “truncated Chl antenna size” in this and other green algae.

FY 2011 Accomplishments

1. Successfully cloned the *Tla2* gene and fully elucidated its function. Manuscript and disclosure in preparation.
2. Successfully cloned the *Tla3* gene. Currently, conducting biochemical analyses and are in the process of elucidating the *Tla3* gene function.

Fiscal Year (FY) 2011 Objectives

- Minimize, or truncate, the chlorophyll antenna size in green algae to maximize photobiological solar conversion efficiency and H₂-production.
- Demonstrate that a truncated chlorophyll (Chl) antenna size would minimize absorption and wasteful dissipation of sunlight by individual cells, resulting in better light utilization efficiency and greater photosynthetic productivity by the green alga mass culture (Table 1).

Technical Barriers

This project addresses the following technical barriers from the Biological Hydrogen Production section of the Fuel Cell Technologies Program Multi-Year Research, Development and Demonstration Plan:

- (AG) Light Utilization Efficiency: Low light utilization efficiency in photobiological hydrogen production due to a large photosystem chlorophyll antenna size.

Technical Targets

The Fuel Cell Technologies Program Multi-Year Plan technical target for 2005 for this project was to reach a 10% utilization efficiency of absorbed light energy (out of a theoretical maximum of 30% possible) in unicellular green. Progress has currently achieved a green alga utilization efficiency of absorbed light energy of about 25% (Table 2).

TABLE 1. *Chlamydomonas reinhardtii* cellular chlorophyll content (Chl/cell), chlorophyll antenna size for photosystem-II (Chl-PSII) and photosystem-I (Chl-PSI), and energy utilization efficiency in wild type, *tla1*, *tla2* and *tla3* mutant strains, as determined by spectrophotometric kinetic analysis (n = 5, ± standard deviation).

	wild type	<i>tla1</i>	<i>tla2</i>	<i>tla3</i>	Long-term goal
Chl/cell mol x10 ⁻¹⁵	2.4 ± 0.5	0.9 ± 0.06	0.93 ± 0.1	0.7 ± 0.1	
Chl-PSII	222 ± 26	115 ± 36	80 ± 30	50 ± 30	37
Chl-PSI	240 ± 4	160 ± 12	115 ± 10	105 ± 10	95
Light Utilization Efficiency (Solar to Chemical)	~3%	~10%	~15%	~25%	~30%

TABLE 2. Progress Achieved vs. the DOE Targets: Utilization Efficiency of Incident Solar Light Energy, E_px E_i

Year	2000	2003	2005	2008	2010	2015
Program Targets	3%	10%*			15%	20%
Actual Progress Achieved	3% Wild Type	10% <i>tla1</i>	15% <i>tla2</i>	25% <i>tla3</i>		

* Target adjusted upward to match ahead-of-schedule progress achieved.



Introduction

The goal of the research is to generate green algal strains with enhanced photosynthetic productivity and H_2 -production under mass culture conditions. To achieve this goal, it is necessary to optimize the light absorption and utilization properties of the cells [1-4]. A cost-effective way to achieve this goal is to reduce the number of Chl molecules that function in the photosystems of photosynthesis. Thus, efforts are under way to isolate microalga mutants with a truncated chlorophyll antenna size.

The rationale for this work is that a truncated light-harvesting Chl antenna size in green algae will prevent individual cells at the surface of the culture from over-absorbing sunlight and wastefully dissipating most of it (Figure 1). A truncated Chl antenna size will permit sunlight to penetrate deeper into the culture, thus enabling many more cells to contribute to useful photosynthesis and H_2 -production (Figure 2). It has been shown that a truncated Chl antenna size will enable about 3-4 times greater solar energy conversion efficiency and photosynthetic productivity than could be achieved with fully pigmented cells [5].

Approach

The focal objective of the research is to identify genes that control the Chl antenna size of photosynthesis and, further, to elucidate how such genes confer a truncated Chl antenna size in the model green alga *Chlamydomonas reinhardtii*. Identification of such genes in *Chlamydomonas* will permit a subsequent transfer of this property, i.e., “truncated Chl antenna size”, to other microalgae of interest




FIGURE 1. Schematic presentation of the fate of absorbed sunlight in fully pigmented (dark green) algae. Individual cells at the surface of the culture over-absorb incoming sunlight (i.e., they absorb more than can be utilized by photosynthesis), and ‘heat dissipate’ most of it. Note that a high probability of absorption by the first layer of cells would cause shading of cells deeper in the culture.

to the DOE Fuel Cell Technologies Program. This objective is currently being approached through DNA insertional mutagenesis/screening and biochemical/molecular/genetic analyses of *Chlamydomonas reinhardtii* cells.

Results

The *tla2* mutant plasmid insert site has been cloned and a gene of interest has been tentatively identified as causing the *tla2* mutation. This molecular and genetic analysis is currently in progress. Work further described the isolation and biochemical and physiological characterization of a new mutant of *Chlamydomonas reinhardtii*, termed *tla3*. Properties of the *tla* “truncated Chl antenna size” strains so far isolated are summarized in Tables 1 and 2, and Figure 3. The *tla3* mutant has the smallest yet Chl antenna size known in green algae.

Future efforts will be directed toward the cloning and characterizing the genes responsible for the *tla* phenotype in *tla2* and *tla3* mutants.

Conclusions

- Significant, ahead-of-schedule progress was achieved in terms of acquiring “truncated Chl antenna size” mutants. This demonstrates feasibility of the approach chosen and success of the methods employed.
- A truncated light-harvesting chlorophyll antenna size in the *tla*-type mutants enhanced solar conversion efficiencies and photosynthetic productivity under bright sunlight conditions.




FIGURE 2. Schematic of sunlight penetration through cells with a truncated chlorophyll antenna size. Individual cells have a diminished probability of absorbing sunlight, thereby permitting penetration of irradiance and H_2 -production by cells deeper in the culture.




FIGURE 3. Project timeline and publications record on the truncated chlorophyll antenna size project. Arrows show publication year of peer-reviewed paper for each of the truncated Chl antenna size mutants. Note that work with the *tla2* and *tla3* strains is now reaching the stage of a peer-reviewed paper.

- Insights on the molecular mechanism for the regulation of the Chl antenna size by the *Tla1* gene were obtained (results not shown pending publication of these findings in a peer reviewed journal).

Future Directions

- Advance the biochemical and molecular characterization of the *tla2* and *tla3* strains.
- Publish results on the *tla2* and *tla3* phenotypes.
- Establish transformation (sense and antisense) protocols with *Tla*-type genes to enhance the down-regulation of the Chl antenna size in *Chlamydomonas reinhardtii*.

FY 2011 Publications/Presentations

1. Peer Reviewed Publication: Ort DR, Zhu X-G, Melis A (2011) Optimizing antenna size to maximize photosynthetic efficiency. *Plant Physiology* 155(1):79-85.
2. DOE Webinar: Melis A (2011) Photosynthesis for Hydrogen and Fuels Production. January 24, 2011. <http://www1.eere.energy.gov/hydrogenandfuelcells/webinar_archives.html>

Patents Issued

1. Melis A and Mitra M (2010) Suppression of *Tla1* gene expression for improved solar energy conversion efficiency and photosynthetic productivity in plants and algae. United States Patent 7,745,696 (issued 29-June-2010).

References

1. Kok B (1953) Experiments on photosynthesis by *Chlorella* in flashing light. In: Burlew JS (ed), *Algal culture: from laboratory to pilot plant*. Carnegie Institution of Washington, Washington D.C., pp 63-75.
2. Myers J (1957) *Algal culture*. In: Kirk RE, Othmer DE (eds), *Encyclopedia of chemical technology*. Interscience, New York, NY, pp 649-668.
3. Radmer R and Kok B (1977) Photosynthesis: Limited yields, unlimited dreams. *Bioscience* 29: 599-605.
4. Mitra M, Melis A (2008) Optical properties of microalgae for enhanced biofuels production. *Optics Express* 16: 21807-21820.
5. Melis A, Neidhardt J and Benemann JR (1999) *Dunaliella salina* (Chlorophyta) with small chlorophyll antenna sizes exhibit higher photosynthetic productivities and photon use efficiencies than normally pigmented cells. *J. appl. Phycol.* 10: 515-525.