

2009 DOE Hydrogen Program

Montana Palladium Research Initiative: Use of Biological Materials and Biologically Inspired Materials for H₂ Catalysis

John W. Peters (PI), Trevor Douglas,
and Mark Young.

Department of Chemistry and Biochemistry
and
Center for Bioinspired Nanomaterials



DOE Project ID#: PDP_19_Peters

This presentation does not contain any proprietary, confidential,
or otherwise restricted information



Overview

Timeline

- Start - Aug. 2006
- End - Dec. 2009

Budget

- Total project funding
 \$1,303,041
 - DOE \$1,031,433

Barriers addressed

- Stability/Durability
- Oxygen Sensitivity
- Electron Donors
- Coupling

Partners

- Montana State University



DOE Project ID#: PDP19





Approaches

Couple Different Catalyst Systems for
Light Driven Hydrogen Generation

Biological catalysts (Hydrogenases)

Nanoparticle biomimetic catalysts




Objectives Addressed


1. Immobilize hydrogenase in gels
2. Determine basis of hydrogenase stability
3. Improve conductivity, mass transfer, and hydrogen production in gels
4. Biomimetic hydrogen production catalyst synthesis
5. Photocatalyst synthesis
6. Coupling catalysts to electrode surfaces
7. H₂ production device fabrication

Approach:

Biological and Biomimetic Catalysts for H₂ production




Hydrogenase Enzymes
(protein architecture protecting
Metal sulfide active site)



Protein encapsulated
nano-catalyst

Coupled Reactions to Generate Hydrogen




GOAL: use **biological catalysts** and develop **biomimetic catalysts** with a variety of sacrificial electron donors or electrochemical source of e^- to produce H_2

Issues and Barriers: Catalyst Stability



- Durability – shelf life
- Reusability
- Product based Inhibition
- Oxygen tolerance / resistance
- Susceptibility to proteolytic inactivation
- Optimization – electron transfer, pH, ionic strength, mediators

Hydrogenases: Highly evolved finely tuned catalysts for *hydrogen oxidation and proton reduction (hydrogen production)*

C. pasteurianum




Desulfovibrio gigas




Cellular location

Membrane Associated
Soluble
Periplasmic
Cytoplasmic

“H Cluster”





NiFe Cluster



Microorganisms:


hydrogen, acetate-grown, methanogenic, green, purple, cyanobacteria; algae; fungus.

Stable NiFe hydrogenase from purple sulfur bacteria forms supermolecular structures




Electron microphotograph of hydrogenase complexes from *T. roseopersicina* negatively stained with 2% uranyl acetate

Cryo reconstruction of hydrogenase from *T. roseopersicina* at ~ 33 Å.




Properties	<i>Thiocapsa roseopersicina</i>
Large subunit	64kDa
Small subunit	34kDa
Temperature optimum	80°C
Stability to Oxygen	stable

Structural studies indicate a role for C-termini in the stability and super molecular complex formation of hydrogenase




B




Sequence alignment of small subunits of stable hydrogenase from *T. roseopersicina* (*T. r.*) and *D. gigas* (*D. g.*) (A) and cryoEM m (B). In the red box the C-termini residues, which could be involve in the cap formation are shown.

With Liang Tang – University of Kansas

Encapsulation of purified active hydrogenases in tetramethyl ortho silicate gels



- Nanoscopic encapsulation;
- Immobilization of unaltered enzyme
- “Heterogeneous material”





Screening electron microscopy photographs of sol-gel

Recovery of hydrogenase activity* encapsulated in Sol-Gel

Hydrogenase	Solution	Gel	Solution/Gel (%)
<i>C. pasterianum</i> (extract)	12550	7581	60.4±16
<i>L. modestogalophilus</i>	9150	6175	67.5±9
<i>T. roseopersicina</i>	12600	8834	70.1±3

*Activity measure at 25° C indicated in nmol/min/mg protein. Values represent average rate over a four-hour period.

Carbon nano tubes incorporated into Sol Gels




Scanning electron microscopy photographs of sol-gel in presence of carbon nanotubes

- Enhance electron transfer
- Facilitate electron transfer between immobilized mediators and hydrogenase
- Facilitate electron transfer between electrodes and hydrogenase in devices


Formation of electro active matrix by encapsulation of the hydrogenase with multiwall carbon nanotubes

Hydrogen production by the silica gel matrix containing hydrogenase (H₂ase), carbon nano tubes (NT) and methyl viologen (MV)





Hydrogenase activity highest in gels with hydrogenase and carbon nanotubes coencapsulated and activity is observed without addition of the redox mediator methyl viologen

The addition of polyethylene glycol (PEG) to the sol gel enhances the hydrogen production by encapsulated hydrogenase




Poly anions and cations modulate the pore size in gels and mass transfer



Controlled synthesis of electroactive polymer gels – controlling protein adsorption, mass transfer, and conductivity







Electrostatic incorporation of protein catalysts




SEM of polymer gels



Biomimetic Catalysts - Synthesis of Pt⁰ Encapsulated Within a Protein Cage Architecture





Coupled Catalysis for H₂ Production



Initial rates (Pt): 4.47x10³ H₂/sec/Hsp
 1.5 x 10⁴ H₂/sec/ferritin
 (Hydrogenase => 6 x 10³ H₂/sec/hydrogenase)

Control of Pt cluster size



(monitored by NCMS) correlation between activity and cluster size
– defining the minimum catalytic cluster



Mass spectra Pt²⁺ bound (black) and Pt⁰ (red) cages - loading of Pt²⁺ (0, 12, 24, 48, 100, and 200 Pt/cage). Charge state 23+ are shown

Moving beyond Pt...

Pd Nanoparticles encapsulated with Ferritin as H₂ Catalysts






Pd particles show significantly lower activity than Pt


Schematic of Pd encapsulated within Ferritin

Stable to 80 °C
Oxygen insensitive



Photocatalyst synthesis - Hematite ($\alpha\text{-Fe}_2\text{O}_3$) in ferritin





$\alpha\text{-Fe}_2\text{O}_3$ is a stable, visible band-gap semiconductor



Attachment of catalysts to electrode surfaces – Cyclic voltammetry to probe e⁻ transfer to catalysts






Protein shell requires an overpotential of ~200mV compared to naked Pt colloid




Attachment of MoS_x - protein cage to GCE - H₂ production

Design and fabrication of prototype devices

Based initially on the solution assay




Testing - Measuring a Photocurrent with $\alpha\text{-Fe}_2\text{O}_3$ Ferritin



Working electrode:
ITO coated glass

Reference electrode:
Ag strip

Counter electrode:
ITO strip



100 mV bias
voltage

Summary

- 1) Hydrogenase can be immobilized in gels and retain activity
- 2) C-termini contributes to stability of stable hydrogenases
- 3) Carbon nanotubes enhance performance of hydrogenase/catalyst in gels
- 4) Pore size in gels can be controlled effecting mass transfer and hydrogenase/catalyst activity
- 5) Pt and Pd hydrogen production catalysts can be synthesized using biological templates
- 6) Photocataysts can be synthesized using biological templates
- 7) Catalysts can be attached to conducting surfaces
- 8) Prototype device has been fabricated and initial testing is underway

Future Work

- 1) Device testing and optimization
- 2) Establish benchmarks for hydrogen production efficiency
- 3) Evaluate hydrogen production efficiency (electrochemical, photochemical, chemical reducing equivalents)
- 4) Evaluate device for durability and sustained H₂ production