

## II.E.1 Maximizing Light Utilization Efficiency and Hydrogen Production in Microalgal Cultures

Tasios Melis

University of California, Berkeley  
 Dept. of Plant & Microbial Biology  
 111 Koshland Hall  
 Berkeley, CA 94720-3102  
 Phone: (510) 642-8166; Fax: (510) 642-4995  
 E-mail: melis@nature.berkeley.edu

DOE Technology Development Manager:

Roxanne Garland

Phone: (202) 586-7260; Fax: (202) 586-9811  
 E-mail: Roxanne.Garland@ee.doe.gov

DOE Project Officer: Carolyn Elam

Phone: (303) 275-4953; Fax: (303) 275-4788  
 E-mail: Carolyn.Elam@go.doe.gov

Contract Number: DE-FG36-05GO15041

Start Date: January 1, 2005

Projected End Date: December 31, 2008

### Objectives

- Minimize, or truncate, the chlorophyll (Chl) antenna size in green algae to maximize photobiological solar conversion efficiency and H<sub>2</sub> production.
- Demonstrate that a truncated Chl antenna size would minimize absorption and wasteful dissipation of sunlight by individual cells, resulting in better light utilization efficiency and greater photosynthetic productivity by the green alga mass culture.

### Technical Barriers

This project addresses the following technical barriers from the Biological Hydrogen Production section (3.1.4.2.5) of the Hydrogen, Fuel Cells and Infrastructure Technologies Program Multi-Year Research, Development and Demonstration Plan:

- (X) Low Light Utilization Efficiency (in Photobiological Hydrogen Production due to a Large Photosystem Chlorophyll Antenna Size)

### Technical Targets

The Hydrogen, Fuel Cells and Infrastructure Technologies Program Multi-Year Research, Development and Demonstration Plan technical target for 2005 for this project was to reach 10% utilization efficiency of absorbed light energy (out of a theoretical maximum of 30% possible) in unicellular green algae.

### Approach

- Employ deoxyribonucleic acid (DNA) insertional mutagenesis, screening, biochemical and molecular genetic analyses for the isolation of “truncated Chl antenna size” mutants in the green alga *Chlamydomonas reinhardtii*.
- Clone and characterize the gene(s) that affect the “Chl antenna size” property in *Chlamydomonas reinhardtii*.
- Apply such genes to generate a “truncated Chl antenna size” in this and other green algae.

### Accomplishments

1. Completed the molecular and genetic analysis of the *tla1* mutant.
2. Completed the functional analysis of the *Tla1* gene.
3. Submitted manuscript on the utility of the *Tla1* gene in conferring a truncated chlorophyll antenna size and on the mechanism by which it maximizes light utilization efficiency and hydrogen production in microalgal cultures.
4. Isolated a new “truncated chlorophyll antenna size” mutant (*tlaX*) with a total of 195 Chl molecules in its photosynthetic apparatus (42% of the wild type). The specific photosystem Chl antenna size measurements of the *tlaX* strain were PSII=80 and PSI=115 Chl molecules.
5. Approximately 15% utilization efficiency of absorbed light energy was achieved (please see Table 1).

**TABLE 1.** Progress Achieved vs. the DOE Targets: Utilization Efficiency of Incident Solar Light Energy, E<sub>0</sub>xE<sub>1</sub>, %.

Year	2000	2003*	2005	2010	2015
Program Targets	3%	10%		15%	20%
Actual Progress Achieved	3%	10%	15%		

\* Target adjusted upward to match ahead-of-schedule progress achieved.

### Introduction

The goal of this project is to generate green algal strains with enhanced photosynthetic productivity and H<sub>2</sub> production under mass culture conditions. To achieve this goal, it is necessary to optimize the light absorption and utilization properties of the cells (Kok


1953; Myers 1957; Radmer and Kok 1977).

A cost-effective way to achieve this goal is to reduce the number of Chl molecules that function in the photosystems of photosynthesis. Thus, efforts are under way to isolate microalga mutants with a truncated chlorophyll antenna size.

The rationale for this R&D is that a truncated light-harvesting Chl antenna size in green algae will prevent individual cells at the surface of the culture from over-absorbing sunlight and wastefully dissipating most of it (Figure 1). A truncated Chl antenna size will permit sunlight to penetrate deeper into the culture, thus enabling many more cells to contribute to useful photosynthesis and H<sub>2</sub> production (Figure 2). It has been shown that a truncated Chl antenna size will enable about 3-4 times greater solar energy conversion efficiency and photosynthetic productivity than could be achieved with fully pigmented cells (Melis et al. 1999).

## Approach

The immediate objective of the research is to identify genes that control the Chl antenna size of photosynthesis and, further, to manipulate such genes so as to confer a truncated Chl antenna size in the model green alga *Chlamydomonas reinhardtii*. Identification of such genes in *Chlamydomonas* will permit a subsequent transfer of this property, i.e., “truncated Chl antenna size”, to other microalgae of interest to the DOE Hydrogen, Fuel Cells and Infrastructure Technologies Program. This objective is currently being approached through DNA insertional mutagenesis/screening and biochemical/molecular/genetic analyses of *Chlamydomonas reinhardtii* cells.




**FIGURE 1.** Schematic Presentation of the Fate of Absorbed Sunlight in Fully Pigmented (Dark Green) Algae. (Individual cells at the surface of the culture over-absorb incoming sunlight [i.e., they absorb more than can be utilized by photosynthesis], and ‘heat dissipate’ most of it. Note that a high probability of absorption by the first layer of cells would cause shading of cells deeper in the culture.)

## Results

The *Chlamydomonas reinhardtii* mutant *tla1* (truncated light-harvesting chlorophyll antenna size) was generated previously upon DNA insertional mutagenesis and shown to specifically possess a smaller than wild type chlorophyll antenna size in both photosystems. This strain exhibited 10% utilization efficiency of incident solar light energy, i.e., substantially greater than the 3% utilization efficiency of the wild type. Molecular and genetic analysis revealed that the exogenous plasmid DNA was inserted at the end of the 5' untranslated region (UTR) and just prior to the Adenine, Thymine, Guanine (ATG) start codon of a hitherto unknown gene (termed *Tla1*), which encodes a protein of 213 amino acids. The *Tla1* gene in the mutant is transcribed with a new 5' UTR sequence, derived from the 3' end of the transforming plasmid. This replacement of the 5' UTR resulted in enhanced transcription of the *tla1* gene in the mutant but inhibition in the translation of the respective *tla1* mRNA. These results provided evidence that down-regulation of the *Tla1* expression is necessary and sufficient to truncate the chlorophyll antenna size and to improve solar utilization efficiencies in a green algal mass culture. Specific applications of the *Tla1* gene in H<sub>2</sub> production were discussed.

Work further described the isolation and biochemical and physiological characterization of a new mutant of *Chlamydomonas reinhardtii*, termed *tlaX*, having a truncated light-harvesting chlorophyll antenna size. This mutant has the smallest Chl antenna size yet known in green algae. This was achieved upon generating and screening an additional 4,500



**FIGURE 2.** Schematic of Sunlight Penetration through Cells with a Truncated Chlorophyll Antenna Size. (Individual cells have a diminished probability of absorbing sunlight, thereby permitting penetration of irradiance and H<sub>2</sub> production by cells deeper in the culture.)

transformants for this purpose, following the protocol of Polle et al. (2003). Properties of the *tlaX* putative “truncated Chl antenna size” strain are summarized in Table 2.

**TABLE 2.** *Chlamydomonas reinhardtii* cellular chlorophyll content, photosystem chlorophyll antenna size and energy utilization efficiency in wild type, *tla1* and *tlaX* mutant strains, as determined by spectrophotometric kinetic analysis (n = 5,  $\pm$ SD).

	wild type	<i>tla1</i>	<i>tlaX</i>	Long-term goal
Chl/cell mol $\times 10^{-15}$	2.4 $\pm$ 0.5	0.9 $\pm$ 0.06	0.93 $\pm$ 0.1	
Chl-PSII	222 $\pm$ 26	115 $\pm$ 36	80 $\pm$ 30	37
Chl-PSI	240 $\pm$ 4	160 $\pm$ 12	115 $\pm$ 10	95
Light Utilization Efficiency	~3%	~10%	~15%	~30%

## Conclusions

- Significant, ahead-of-schedule progress was achieved in terms of acquiring “truncated Chl antenna size” mutants. *This demonstrates feasibility of the approach chosen and success of the methods employed.*
- A truncated light-harvesting chlorophyll antenna size in the *tla*-type mutants leads to enhanced solar conversion efficiencies and greater photosynthetic productivity of the algae under bright sunlight conditions.
- Insights on the molecular mechanism for the regulation of the Chl antenna size by the *Tla1* gene were obtained (results not shown pending publication of these findings in a peer reviewed journal).

## Future Directions

- Advance the biochemical and molecular characterization of the *tlaX* strain.
- Establish transformation (sense and antisense) protocols with *Tla*-type genes to enhance the down-regulation of the Chl antenna size in *Chlamydomonas reinhardtii*.
- Perform comparative green-alga light utilization efficiency and photosynthetic productivity measurements under mass culture conditions in wild type and *tla*-type mutants.
- Isolate additional *tla*-type mutants.
- Perform genetic crosses to combine different *tla*-type properties.

## References

1. Kok B. (1953) Experiments on photosynthesis by *Chlorella* in flashing light. In: Burlew J.S. (ed), *Algal culture: from laboratory to pilot plant*. Carnegie Institution of Washington, Washington D.C., pp 63-75.
2. Melis A., Neidhardt J. and Benemann J.R. (1999) *Dunaliella salina* (Chlorophyta) with small chlorophyll antenna sizes exhibit higher photosynthetic productivities and photon use efficiencies than normally pigmented cells. *J. appl. Phycol.* 10: 515-52.
3. Myers J. (1957) *Algal culture*. In: Kirk R.E., Othmer D.E. (eds), *Encyclopedia of chemical technology*. Interscience, New York, NY, pp. 649-668.
4. Polle J.E.W., Kanakagiri S. and Melis A. (2003) *tla1*, a DNA insertional transformant of the green alga *Chlamydomonas reinhardtii* with a truncated light-harvesting chlorophyll antenna size. *Planta* 217: 49-59.
5. Radmer R. and Kok B. (1977) Photosynthesis: Limited yields, unlimited dreams. *Bioscience* 29: 599-605.

## FY 2006 Publications/Presentations

1. Melis A. (2005) Bioengineering of green algae to enhance photosynthesis and hydrogen production. Chapter 12 in *Artificial Photosynthesis: From Basic Biology to Industrial Application*, A.F. Collins and C. Critchley (eds.), Wiley-Verlag & Co., pp. 229-240.
2. White A.L. and Melis, A. (2006) Biochemistry of hydrogen metabolism in *Chlamydomonas reinhardtii* wild type and a Rubisco-less mutant. *Intl. J. Hydrogen Energy* 31: 455-464.
3. Melis A., Zhang L., Benemann J.R., Forestier M., Ghirardi M.L., Seibert M. (2006) Hydrogen production using hydrogenase-containing oxygenic photosynthetic organisms. United States Patent 6,989,252 B2 (issued 24-Jan-2006).
4. Tetali S.D., Mitra M., Melis A. (2006) *Tla1*, a nuclear gene regulates the light-harvesting chlorophyll antenna size in the green alga *Chlamydomonas reinhardtii*. (Submitted for publication).
5. Mitra M., Kanakagiri, S.D. and Melis A. (2005) Chlorophyll antenna size adjustments in *Chlamydomonas reinhardtii* involve coordinate regulation of *Tla1*, *CAO* and *Lhcb* gene expression. *Plant Biology* 2005, Abstract #236 in the American Society of Plant Biologists 2005 Annual Meeting Program Book. p. 142.
6. Melis A. (2005) Integrated photobiological H-2 production. Society for Industrial Microbiology and Biotechnology (SIM) Annual Meeting Program and Abstracts. Abstract S43, p. 65.
7. Mitra M., Kanakagiri S.D. and Melis A. (2006) Chlorophyll antenna size adjustments by irradiance in *Chlamydomonas reinhardtii* involve coordinate regulation of *Tla1*, *CAO* and *Lhcb* gene expression. 15<sup>th</sup> Western Photosynthesis Conference 5-8 January 2006 book of abstracts. p. 22-23.